ELSEVIER

Environmental Pollution 136 (2005) I–IV

ENVIRONMENTAL POLLUTION

www.elsevier.com/locate/envpol

Contents of Volume 136

Number 1 2005

1 **Seasonal and intertidal impact on DNA adduct levels in gills of blue mussels (*Mytilus edulis* L.)**
H. Skarphéðinsdóttir, G. Ericson, H.P. Halldórsson, J. Svavarsson

Gill DNA adduct and total PAH tissue levels were higher in mussels in the intertidal than subtidal zone, and higher in winter than summer.

11 **Potentially toxic metals in ombrotrophic peat along a 400 km English–Scottish transect**
E.J. Smith, S. Hughes, A.J. Lawlor, S. Lofts, B.M. Simon, P.A. Stevens, R.T. Stidson, E. Tipping, C.D. Vincent

Peat metal contents depend upon proximity to industrial and mining areas; the metals may be exerting toxic effects in some places.

19 **Flame retardants, surfactants and organotins in sediment and mysid shrimp of the Scheldt estuary (The Netherlands)**
T.A. Verslycke, A.D. Vethaak, K. Arijs, C.R. Janssen

Field populations of mysid shrimp (Neomysis integer) of the Scheldt estuary (The Netherlands) are exposed to high concentrations of endocrine disruptors.

33 **Seasonal trends in reduced leaf gas exchange and ozone-induced foliar injury in three ozone sensitive woody plant species**
K. Novak, M. Schaub, J. Fuhrer, J.M. Skelly, C. Hug, W. Landolt, P. Bleuler, N. Kräuchi

Reductions in leaf gas exchange corresponded to the onset of ozone-induced visible foliar injury for seedlings exposed to ambient ozone exposures.

47 **Lead and cadmium in indoor air and the urban environment**
G.J.K. Komarnicki

Urban vegetation and isopods are potential indicators for indoor aerial heavy metals.

63 **A novel modelling approach for spatial and temporal variations in nitrate concentrations in an N-impacted UK small upland river basin**
R.P. Smart, M.S. Cresser, L.J. Calver, M. Clark, P.J. Chapman

A novel model of spatial and temporal variations in stream water nitrate concentration throughout an N-impacted upland catchment is described.

71 **Serum concentrations and profiles of polychlorinated biphenyls in Taiwan Yu-cheng victims twenty years after the incident**
S.-C.C. Lung, Y.-L.L. Guo, H.-Y. Chang

Total serum PCBs in the Yu-cheng adult victims twenty years after the incident were still higher than that of the general population in Taiwan.

81 **Brominated flame retardants and organochlorine pollutants in eggs of little owls (*Athene noctua*) from Belgium**
V. Jaspers, A. Covaci, J. Maervoet, T. Dauwe, S. Voorspoels, P. Schepens, M. Eens

PBDEs are measurable in Belgian little owl eggs, but at relatively low concentrations compared to other birds of prey.

89 **Long-term trends and spatial variability in nitrate leaching from alpine catchment–lake ecosystems in the Tatra Mountains (Slovakia–Poland)**
J. Kopáček, E. Stuchlík, R.F. Wright

An empirical model explains 80% of the observed spatial and temporal variability in the long-term (1937–2000) alpine lake water nitrate concentrations in the Tatra Mountains.

103 **Stress biomarkers and alkali-labile phosphate level in mussels (*Mytilus galloprovincialis*) collected in the urban area of Venice (Venice Lagoon, Italy)**
D.M. Pampanin, I. Marangon, E. Volpato, G. Campesan, C. Nasci

The overall biological and chemical data show a higher stress condition in the mussels (Mytilus galloprovincialis) from the urban area of Venice.

109 **Vegetation composition of roadside verges in Scotland: the effects of nitrogen deposition, disturbance and management**
A.M. Truscott, S.C.F. Palmer, G.M. McGowan, J.N. Cape, S. Smart

NO_x, NH_3 and road verge vegetation Ellenberg fertility indices decline with distance from traffic.

S0269–7491(05)00184-3

119 Trace elements in biodeposits and sediments from mussel culture in the Ría de Arousa (Galicia, NW Spain)
X.L. Otero, P. Vidal-Torrado, R.M. Calvo de Anta, F. Macías

Geochemical behaviour of trace metals in mussel biodeposits and sediments were studied.

135 Copper regulation and homeostasis of *Daphnia magna* and *Pseudokirchneriella subcapitata*: influence of acclimation
B.T.A. Bossuyt, C.R. Janssen

A freshwater alga and zooplankter maintained copper homeostasis, particularly when acclimated to high copper levels.

145 Copper toxicity to different field-collected cladoceran species: intra- and inter-species sensitivity
B.T.A. Bossuyt, C.R. Janssen

Zooplankton living in natural waters are more sensitive to copper than laboratory animals.

155 Human health risk assessment of organochlorines associated with fish consumption in a coastal city in China
Q.T. Jiang, T.K.M. Lee, K. Chen, H.L. Wong, J.S. Zheng, J.P. Giesy, K.K.W. Lo, N. Yamashita, P.K.S. Lam

Health risk assessment of organochlorines associated with fish consumption reveals potential cancer risks for some contaminants in a coastal population in China.

167 Thermal conductivity of leaf compost used in biofilters: An experimental and theoretical investigation
M. Chandrakanthi, A.K. Mehrotra, J.P.A. Hettiaratchi

Thermal conductivity of compost exhibits a linear relationship with the degree of saturation of the matrix.

175 Relationships between ecological variables and four organochlorine pollutants in an artic glaucous gull (*Larus hyperboreus*) population
J.O. Bustnes, Ø. Miland, M. Fjeld, K.E. Erikstad, J.U. Skaare

In arctic glaucous gulls, nesting behaviour, early chick growth and adult return rate were negatively related to blood concentration of organochlorines.

187 Bioremediation of petroleum hydrocarbons in contaminated soils: Comparison of biosolids addition, carbon supplementation, and monitored natural attenuation
D. Sarkar, M. Ferguson, R. Datta, S. Birnbaum

Addition of biosolids is a potentially effective method of biostimulation for degradation of petroleum hydrocarbons in soils.

Number 2 2005

197 Bioassay responses and effects on benthos after pilot remediations in the delta of the rivers Rhine and Meuse
P.J. den Besten, P.J. van den Brink

Capping contaminated sediments can be an effective remediation measure in two large river deltas.

209 Identifying the source, transport path and sinks of sewage derived organic matter
S.M. Mudge, C.E. Duce

Multivariate statistical analysis was used to identify pathways and accumulation sites for contaminants in coastal waters.

221 Identification of a new Irgarol-1051 related *s*-triazine species in coastal waters
K.-H. Lam, Z. Cai, H.-Y. Wai, V.-H. Tsang, M.-W. Lam, R.-H. Cheung, H. Yu, P.-S. Lam

An s-triazine species resists degradation and may be a chemical risk for marine coastal communities.

231 Cytochrome P4501A induction in brown trout exposed to small streams of an urbanised area: results of a five-year-study
A. Behrens, H. Segner

The CYP1A biomarker in brown trout robustly ranks the chemical stress status of small streams.

243 The reproductive success and quality of blue tits (*Parus caeruleus*) in a heavy metal pollution gradient
T. Dauwe, E. Janssens, R. Pinxten, M. Eens

Exposure to elevated heavy metal concentrations did not result in a significant reduction of the reproductive success in free-living blue tits.

253 Organochlorine compounds in soils and sediments of the mountain Andean Lakes
F. Borghini, J.O. Grimalt, J.C. Sanchez-Hernandez, R. Barra, C.J.T. García, S. Focardi

Distribution of persistent organochlorine compounds in the Andean Mountain lakes is influenced by temperature.

267 How to accurately assay the algal toxicity of pesticides with low water solubility
J. Ma, J. Chen

A new method is described for assaying the toxicity of water insoluble pesticides.

275 Influence of soil zinc concentrations on zinc sensitivity and functional diversity of microbial communities
K. Lock, C.R. Janssen

Zinc sensitivity of microbial communities and functional diversity decrease with increasing zinc concentration in the pore water.

283 **Temporal and spatial variation of perchlorate in streambed sediments: results from in-situ dialysis samplers**
K. Tan, T.A. Anderson, W.A. Jackson

Perchlorate may be rapidly attenuated in saturated near-surface sediments.

293 **Transplants of aquatic mosses as biomonitors of metals released by a mine effluent**
R. Figueira, T. Ribeiro

Effects of metal contamination could be related to changes in intracellular concentrations of Mg.

303 **Global pollution monitoring of polychlorinated dibenzo-*p*-dioxins (PCDDs), furans (PCDFs) and coplanar polychlorinated biphenyls (coplanar PCBs) using skipjack tuna as bioindicator**
D. Ueno, M. Watanabe, A. Subramanian, H. Tanaka, G. Fillmann, P.K.S. Lam, G.J. Zheng, M. Muchtar, H. Razak, M. Prudente, K.-H. Chung, S. Tanabe

Global pollution monitoring of PCDD/Fs and coplanar PCBs in offshore water and open sea were conducted using skipjack tuna as bioindicator.

315 **Lead and zinc bioavailability to *Eisenia fetida* after phosphorus amendment to repository soils**
D.R. Ownby, K.A. Galvan, M.J. Lydy

Phosphorus form and pH were controlling factors in the effectiveness of phosphorus amendment in decreasing Pb and Zn bioavailability.

323 **Preliminary screening of perfluorooctane sulfonate (PFOS) and other fluorochemicals in fish, birds and marine mammals from Greenland and the Faroe Islands**
R. Bossi, F.F. Riget, R. Dietz, C. Sonne, P. Fauser, M. Dam, K. Vorkamp

Perfluorinated acids were detected in livers of fish, birds and marine mammals from Greenland and the Faroe Islands.

331 **The impact of the Almalyk Industrial Complex on soil chemical and biological properties**
N. Shukurov, S. Pen-Mouratov, Y. Steinberger

Soil nematodes and microbes are suitable biomonitors for metals in soils.

341 **Mercury volatilisation and phytoextraction from base-metal mine tailings**
F.N. Moreno, C.W.N. Anderson, R.B. Stewart, B.H. Robinson

Mass balance studies indicated that volatilisation was a dominant pathway for mercury removal.

353 **Species- and stage-specific differences in trace element tissue concentrations in amphibians: implications for the disposal of coal-combustion wastes**
J.H. Roe, W.A. Hopkins, B.P. Jackson

Results suggest that metamorphosed amphibians can transport trace elements from aquatic disposal basins to non-contaminated habitats.

Number 3 2005

New initiative

365 **Response patterns in adult forest trees to chronic ozone stress: identification of variations and consistencies**
A.J. Nunn, I.M. Reiter, K.-H. Häberle, C. Langebartels, G. Bahnweg, H. Pretzsch, H. Sandermann, R. Matyssek

Responses of adult forest trees to chronic O_3 stress can be visualized in a survey table applying a three-grade colour coding to each investigated parameter.

Regular papers

371 **Responses of sensitive and tolerant bush beans (*Phaseolus vulgaris* L.) to ozone in open-top chambers are influenced by phenotypic differences, morphological characteristics, and the chamber environment**
V. Elagöz, W.J. Manning

Phenotypic differences, morphological characteristics, and 'chamber effects' inside OTCs are equally influential in determining the responses of beans to O_3.

385 **Factors affecting the effects of EDU on growth and yield of field-grown bush beans (*Phaseolus vulgaris* L.), with varying degrees of sensitivity to ozone**
V. Elagöz, W.J. Manning

Plant sensitivity to ozone, stage of plant development, number of applications of EDU and ambient ozone affect bean plant responses to EDU.

397 **Accumulating pollutants in conifer needles on an Atlantic island – A case study with on Tenerife, Canary Islands**
M. Tausz, W. Trummer, W. Goessler, A. Wonisch, D. Grill, S. Naumann, M.S. Jiménez, D. Morales

Pollutant concentrations in pine needles indicate moderate local impact of sulphur, particulate matter, and sea sprays at lower elevation plots in Tenerife.

409 Bioavailability of contaminants estimated from uptake rates into soil invertebrates
N.M. van Straalen, M.H. Donker, M.G. Vijver, C.A.M. van Gestel

Sublethal toxicity of zinc to isopods suggests that bioavailability of soil contaminants is best measured by uptake rates, not by body burdens.

419 Selected chlorobornanes, polychlorinated naphthalenes and brominated flame retardants in Bjørnøya (Bear Island) freshwater biota
A. Evenset, G.N. Christensen, R. Kallenborn

Elevate levels of chlorobornanes, polychlorinated naphthalenes and brominated flame retardants have been measured in biota from a Norwegian Arctic lake.

431 Atmospheric trace metal pollution in the Naples urban area based on results from moss and lichen bags
S. Giordano, P. Adamo, S. Sorbo, S. Vingiani

Despite significant damage to tissue and cell integrity, moss and lichen in bags efficiently intercept airborne trace elements.

443 Earthworm responses to Cd and Cu under fluctuating environmental conditions: A comparison with results from laboratory exposures
D.J. Spurgeon, C. Svendsen, L.J. Lister, P.K. Hankard, P. Kille

Comparative analysis of life-cycle, physiological and molecular responses to Cu and Cd indicate similar responses under static and fluctuating climate regimes.

453 Uptake, absorption efficiency and elimination of DDT in marine phytoplankton, copepods and fish
X. Wang, W.-X. Wang

Biomagnification and exposure of DDT in a marine food chain is demonstrated by measurements of uptake and elimination rates and kinetic modeling.

465 Contamination status and accumulation features of persistent organochiorines in pet dogs and cats from Japan
T. Kunisue, S. Nakanishi, M. Watanabe, T. Abe, S. Nakatsu, S. Kawauchi, A. Sano, A. Horii, Y. Kano, S. Tanabe

Pet dogs may have higher metabolic and elimination capacity for organochlorines than pet cats.

477 Benzo[*a*]pyrene co-metabolism in the presence of plant root extracts and exudates: Implications for phytoremediation
J.A. Rentz, P.J.J. Alvarez, J.L. Schnoor

Bacterial benzo[a]pyrene cometabolism, a plant-microbe interaction affecting polycyclic aromatic hydrocarbon phytoremediation was demonstrated with Sphingomonas yanoikuyae JAR02 that utilized plant root extracts and exudates as primary substrates.

485 Metal-contaminated soil remediation by means of paper mill sludges addition: chemical and ecotoxicological evaluation
N. Calace, T. Campisi, A. Iacondini, M. Leoni, B.M. Petronio, M. Pietroletti

Paper mill sludge decreased available metals.

493 Occurrence of polybrominated diphenylethers, polychlorinated dibenzo-*p*-dioxins, dibenzofurans and biphenyls in coastal sediments from Spain
E. Eljarrat, A. De La Cal, D. Larrazabal, B. Fabrellas, A.R. Fernandez-Alba, F. Borrull, R.M. Marce, D. Barcelo

Decabrominated diphenyl ether makes a substantial contribution to the total PBDE contamination found in Spanish coastal sediments.

503 Review of passive accumulation devices for monitoring organic micropollutants in the aquatic environment
F. Stuer-Lauridsen

Major developments in the passive sampling of organic contaminants in aquatic environments will support future monitoring, compliance and research.

525 Survey of organotin compounds in rivers and coastal environments in Portugal 1999–2000
S. Díez, S. Lacorte, P. Viana, D. Barceló, J.M. Bayona

Survey data illustrate the complexities of organotin sources and cycling in coastal environments.